MILWAUKEE PUBLIC SCHOOLS, WI

MIDDLE SCHOOL YOUTH RISK BEHAVIOR SURVEY
2003

Table 2k. Percentage of Middle School Students Who Had Rarely or Never Worn Seatbelts* and Who Had Ever Ridden in a Car with a Driver Who Had Been Drinking Alcohol, by Sex, Race/ Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | M | lale | Female | | Total | |
|-----------------------------------|-------------|---------------------|--------|--------|-------|--------|
| Question | % | CI [†] (±) | % | CI (±) | % | CI (±) |
| Rarely or never wore seat belts | | | | | | |
| Race/Ethnicity | | | | | | |
| White [§] | 24.3 | 6.8 | 15.1 | 9.0 | 20.4 | 6.4 |
| Black§ | 35.7 | 6.1 | 26.0 | 4.4 | 30.9 | 4.2 |
| Hispanic | 31.5 | 6.8 | 19.6 | 7.1 | 25.7 | 4.9 |
| Other | _1 | _ | _ | _ | 25.1 | 7.6 |
| Grade | | | | | | |
| 6 | 27.6 | 7.2 | 13.9 | 5.1 | 20.9 | 4.6 |
| 7 | 30.8 | 4.9 | 28.1 | 3.9 | 29.4 | 3.3 |
| 8 | 39.4 | 7.4 | 26.7 | 6.8 | 33.1 | 5.8 |
| Total | 32.8 | 4.1 | 23.3 | 4.0 | 28.1 | 3.3 |
| Rode with a driver who had been d | rinking alc | ohol | | | | |
| Race/Ethnicity | | | | | | |
| White | 38.3 | 6.9 | 43.7 | 7.8 | 40.7 | 5.2 |
| Black | 40.6 | 5.6 | 37.9 | 5.3 | 39.2 | 3.6 |
| Hispanic | 33.6 | 10.1 | 37.1 | 8.7 | 35.4 | 8.0 |
| Other | _ | _ | _ | _ | 37.0 | 8.0 |
| Grade | | | | | | |
| 6 | 35.9 | 10.9 | 35.7 | 5.8 | 35.7 | 7.2 |
| 7 | 36.0 | 4.7 | 36.5 | 7.8 | 36.0 | 4.9 |
| 8 | 42.5 | 4.8 | 43.5 | 6.9 | 43.0 | 4.5 |
| Total | 38.3 | 4.3 | 38.8 | 4.0 | 38.5 | 3.1 |

^{*}When riding in a car.

^{†95%} confidence interval.

[§]Non-Hispanic.

¹Not available.

Table 3k. Percentage of Middle School Students Who Had Rarely or Never Worn Bicycle Helmets* or Rollerblade or Skateboard Helmets,† by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | M | lale | Female | | Total | |
|---------------------------|---------------------|---------------------|---------------|--------|-------|--------|
| Question | % | CI [§] (±) | % | CI (±) | % | CI (±) |
| Rarely or never wore bicy | cle helmets | | | | | |
| Race/Ethnicity | | | | | | |
| White ¹ | 83.7 | 12.1 | ** | _ | 82.2 | 7.9 |
| Black ¹ | 95.7 | 3.0 | 93.9 | 3.0 | 94.8 | 2.6 |
| Hispanic | 91.0 | 4.3 | 85.2 | 6.3 | 88.3 | 4.3 |
| Other | _ | _ | _ | _ | 87.9 | 4.4 |
| Grade | | | | | | |
| 6 | 90.4 | 7.8 | 83.2 | 4.9 | 87.0 | 5.0 |
| 7 | 93.7 | 3.2 | 92.2 | 3.5 | 93.0 | 2.5 |
| 8 | 93.9 | 2.9 | 94.5 | 3.5 | 94.2 | 2.9 |
| Total | 92.6 | 2.9 | 90.1 | 2.6 | 91.4 | 2.1 |
| Rarely or never wore a he | lmet when using rol | lerblades or ridi | ng a skateboa | rd | | |
| Race/Ethnicity | | | | | | |
| White | _ | _ | _ | _ | 84.5 | 6.7 |
| Black | 88.3 | 5.3 | 94.4 | 3.2 | 91.6 | 3.3 |
| Hispanic | 84.4 | 8.4 | 86.7 | 6.4 | 85.2 | 4.2 |
| Other | _ | _ | _ | _ | 90.5 | 6.2 |
| Grade | | | | | | |
| 6 | 84.4 | 8.3 | 86.1 | 7.1 | 85.2 | 4.9 |
| 7 | 85.3 | 6.7 | 92.5 | 4.1 | 89.1 | 4.2 |
| 8 | 90.8 | 3.5 | 94.9 | 3.2 | 92.9 | 2.8 |
| Total | 86.6 | 4.0 | 91.0 | 2.7 | 88.8 | 2.3 |

^{*}Among the 90.6% of students who had ridden a bicycle.

 $^{^{\}scriptscriptstyle \dagger}\!A\text{mong}$ the 56.8% of students who used rollerblades or ridden a skateboard.

^{§95%} confidence interval.

¹Non-Hispanic.

^{**}Not available.

Table 4k. Percentage of Middle School Students Who Had Ever Carried a Weapon,* Who Had Ever Been in a Physical Fight, and Who Had Ever Been Hurt in a Physical Fight,† by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | M | lale | Female | | Total | |
|-------------------------------|------|---------|--------|--------|-------|--------|
| Question | % | CI§ (±) | % | CI (±) | % | CI (±) |
| Carried a weapon | | | | | | |
| Race/Ethnicity | | | | | | |
| White ¹ | 35.4 | 8.8 | 20.7 | 9.0 | 29.0 | 5.7 |
| Black ¹ | 36.5 | 4.4 | 24.2 | 4.5 | 30.3 | 3.5 |
| Hispanic | 31.8 | 11.5 | 23.9 | 8.0 | 27.9 | 7.8 |
| Other | ** | _ | _ | _ | 31.2 | 6.8 |
| Grade | | | | | | |
| 6 | 31.2 | 9.1 | 18.7 | 5.0 | 25.1 | 5.5 |
| 7 | 29.3 | 5.7 | 23.4 | 4.9 | 26.4 | 3.4 |
| 8 | 45.4 | 4.6 | 28.8 | 6.2 | 37.2 | 3.7 |
| Total | 35.3 | 4.2 | 23.9 | 3.3 | 29.7 | 2.9 |
| Were in a physical fight | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 75.1 | 6.3 | 45.9 | 10.5 | 62.6 | 7.7 |
| Black | 78.8 | 5.1 | 75.1 | 4.8 | 77.0 | 3.8 |
| Hispanic | 74.9 | 7.0 | 58.1 | 9.5 | 66.7 | 7.3 |
| Other | _ | _ | _ | _ | 72.8 | 7.0 |
| Grade | | | | | | |
| 6 | 71.1 | 8.1 | 59.8 | 11.0 | 65.6 | 7.7 |
| 7 | 75.6 | 5.1 | 68.6 | 5.7 | 72.3 | 4.2 |
| 8 | 84.7 | 4.6 | 75.9 | 5.6 | 80.4 | 3.1 |
| Total | 77.2 | 3.7 | 68.3 | 4.6 | 72.9 | 3.3 |
| Were hurt in a physical fight | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 11.2 | 4.5 | 6.3 | 4.8 | 9.1 | 3.1 |
| Black | 13.1 | 4.6 | 6.6 | 2.6 | 9.8 | 2.8 |
| Hispanic | 11.9 | 5.6 | 7.0 | 3.5 | 9.4 | 3.9 |
| Other | _ | _ | _ | _ | 6.1 | 4.6 |
| Grade | | | | | | |
| 6 | 16.6 | 7.9 | 5.3 | 4.1 | 11.1 | 4.7 |
| 7 | 9.4 | 2.5 | 7.4 | 3.3 | 8.3 | 2.3 |
| 8 | 11.1 | 3.8 | 7.3 | 3.1 | 9.3 | 2.7 |
| Total | 12.2 | 3.1 | 6.7 | 2.0 | 9.5 | 2.0 |

^{*}For example, a gun, knife, or club.

[†]Had to be treated by a doctor or nurse.

^{§95%} confidence interval.

¹Non-Hispanic.

^{**}Not available.

Table 5k. Percentage of Middle School Students Who Had Ever Seriously Thought About Killing Themselves, Who Had Ever Made a Plan About How To Kill Themselves, and Who Had Ever Tried To Kill Themselves, by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | M | Male | | Female | | Total | |
|-----------------------------|-----------------|---------|------|--------|------|--------|--|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) | |
| Seriously thought about kil | ling themselves | | | | | | |
| Race/Ethnicity | | | | | | | |
| White [†] | 21.7 | 7.3 | 31.6 | 9.6 | 25.9 | 5.8 | |
| Black [†] | 19.8 | 3.8 | 27.5 | 4.6 | 23.6 | 3.1 | |
| Hispanic | 15.1 | 8.0 | 33.2 | 10.9 | 24.1 | 8.7 | |
| Other | § | _ | _ | _ | 24.8 | 6.9 | |
| Grade | | | | | | | |
| 6 | 18.2 | 6.4 | 24.1 | 9.5 | 21.0 | 6.4 | |
| 7 | 15.8 | 4.8 | 27.8 | 5.8 | 21.5 | 4.2 | |
| 8 | 24.0 | 4.5 | 34.9 | 5.7 | 29.3 | 3.4 | |
| Total | 19.2 | 2.8 | 29.0 | 4.2 | 24.0 | 2.9 | |
| Made a plan about how to l | kill themselves | | | | | | |
| Race/Ethnicity | | | | | | | |
| White | 7.7 | 5.6 | 17.0 | 10.4 | 11.7 | 6.1 | |
| Black | 8.1 | 2.5 | 16.2 | 3.4 | 12.1 | 2.4 | |
| Hispanic | 10.3 | 5.6 | 15.6 | 7.5 | 13.2 | 5.9 | |
| Other | _ | _ | _ | _ | 18.6 | 6.6 | |
| Grade | | | | | | | |
| 6 | 6.1 | 3.1 | 13.3 | 5.5 | 9.6 | 3.4 | |
| 7 | 9.2 | 3.6 | 15.3 | 4.2 | 12.1 | 3.3 | |
| 8 | 10.4 | 3.4 | 21.4 | 5.5 | 15.8 | 3.3 | |
| Total | 8.7 | 2.1 | 16.7 | 3.2 | 12.6 | 2.2 | |
| Tried to kill themselves | | | | | | | |
| Race/Ethnicity | | | | | | | |
| White | 5.9 | 4.4 | 14.5 | 7.8 | 9.6 | 4.6 | |
| Black | 8.7 | 3.4 | 13.1 | 3.5 | 10.9 | 2.6 | |
| Hispanic | 6.8 | 3.7 | 11.2 | 4.3 | 9.0 | 3.4 | |
| Other | _ | _ | _ | _ | 15.5 | 7.0 | |
| Grade | | | | | | | |
| 6 | 7.1 | 4.3 | 10.9 | 5.2 | 9.0 | 3.6 | |
| 7 | 7.8 | 4.1 | 12.9 | 3.5 | 10.2 | 3.2 | |
| 8 | 8.5 | 2.8 | 16.3 | 4.2 | 12.3 | 2.6 | |
| Total | 8.1 | 2.3 | 13.4 | 2.6 | 10.6 | 2.0 | |

^{*95%} confidence interval.

[†]Non-Hispanic.

[§]Not available.

Table 6k. Percentage of Middle School Students Who Had Ever Smoked Cigarettes, by Sex, Race/ Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | M | lale | Female | | Total | |
|-------------------------------------|----------|---------|--------|--------|-------|--------|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) |
| Lifetime cigarette use [†] | | | | | | |
| Race/Ethnicity | | | | | | |
| White§ | 32.5 | 12.7 | 36.3 | 8.3 | 34.2 | 8.4 |
| Black [§] | 49.8 | 4.9 | 52.7 | 4.7 | 51.2 | 3.6 |
| Hispanic | 35.4 | 9.2 | 42.4 | 12.1 | 38.9 | 9.2 |
| Other | 1 | _ | _ | _ | 43.0 | 7.1 |
| Grade | | | | | | |
| 6 | 34.5 | 10.7 | 37.0 | 10.2 | 35.6 | 8.1 |
| 7 | 40.3 | 6.0 | 51.5 | 6.3 | 45.8 | 5.2 |
| 8 | 56.3 | 5.2 | 56.4 | 6.0 | 56.4 | 4.4 |
| Total | 43.9 | 4.7 | 48.6 | 4.3 | 46.2 | 3.7 |
| Lifetime daily cigarette use** | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 3.4 | 3.6 | 10.6 | 7.4 | 6.5 | 3.4 |
| Black | 5.4 | 1.9 | 4.0 | 1.7 | 4.7 | 1.6 |
| Hispanic | 6.1 | 4.6 | 4.6 | 3.6 | 5.3 | 2.9 |
| Other | _ | _ | _ | _ | 8.4 | 5.1 |
| Grade | | | | | | |
| 6 | 2.7 | 2.5 | 4.8 | 3.9 | 3.7 | 2.3 |
| 7 | 7.5 | 3.2 | 4.4 | 2.1 | 5.9 | 2.4 |
| 8 | 5.3 | 2.6 | 6.5 | 2.8 | 5.9 | 1.7 |
| Total | 5.2 | 1.6 | 5.3 | 1.5 | 5.3 | 1.2 |

^{*95%} confidence interval.

[†]Ever tried cigarette smoking, even one or two puffs.

[§]Non-Hispanic.

¹Not available.

^{**}Ever smoked at least one cigarette every day for 30 days.

Table 7k. Percentage of Middle School Students Who Smoked Cigarettes, by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | 1 | 1ale | Female | | Total | |
|------------------------------------|-----|---------|--------|--------|-------|--------|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) |
| Current cigarette use [†] | | | | | | |
| Race/Ethnicity | | | | | | |
| White [§] | 7.0 | 5.3 | 11.8 | 4.6 | 9.1 | 3.6 |
| Black§ | 5.9 | 2.1 | 7.2 | 2.2 | 6.5 | 1.5 |
| Hispanic | 5.8 | 3.6 | 11.2 | 5.0 | 8.7 | 3.1 |
| Other | _1 | _ | _ | _ | 10.0 | 5.2 |
| Grade | | | | | | |
| 6 | 4.1 | 3.4 | 6.5 | 4.4 | 5.2 | 2.7 |
| 7 | 5.9 | 3.0 | 8.3 | 3.2 | 7.0 | 2.4 |
| 8 | 8.7 | 3.7 | 10.7 | 3.4 | 9.7 | 2.4 |
| Total | 6.2 | 1.8 | 8.6 | 1.7 | 7.4 | 1.2 |
| Current frequent cigarette use** | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 1.4 | 2.1 | 1.8 | 2.2 | 1.6 | 1.6 |
| Black | 1.4 | 1.0 | 0.8 | 0.7 | 1.1 | 0.6 |
| Hispanic | 0.7 | 1.2 | 0.9 | 1.2 | 0.8 | 0.9 |
| Other | _ | _ | _ | _ | 3.6 | 3.0 |
| Grade | | | | | | |
| 6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7 | 2.1 | 2.2 | 0.7 | 0.9 | 1.4 | 1.2 |
| 8 | 1.8 | 1.4 | 2.3 | 1.6 | 2.1 | 1.1 |
| Total | 1.3 | 0.8 | 1.1 | 0.7 | 1.3 | 0.5 |

^{*95%} confidence interval.

 $^{^{\}dagger}\text{Smoked}$ cigarettes on ≥ 1 of the 30 days preceding the survey.

[§]Non-Hispanic.

¹Not available.

^{**}Smoked cigarettes on \geq 20 of the 30 days preceding the survey.

Table 8k. Percentage of Middle School Students Who Used Smokeless Tobacco,* Smoked Cigars,† and Used Any Tobacco Product,§ by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | M | 1ale | Female | | Total | |
|-------------------------------|------|---------------------|--------|--------|-------|--------|
| Question | % | CI ¹ (±) | % | CI (±) | % | CI (±) |
| Current smokeless tobacco use | | | | | | |
| Race/Ethnicity | | | | | | |
| White** | 1.4 | 2.0 | 3.1 | 3.4 | 2.1 | 1.9 |
| Black** | 2.6 | 1.7 | 1.7 | 1.2 | 2.1 | 1.1 |
| Hispanic | 6.5 | 5.1 | 3.9 | 3.4 | 5.2 | 2.7 |
| Other | †† | _ | _ | _ | 0.9 | 1.0 |
| Grade | | | | | | |
| 6 | 2.5 | 2.8 | 2.6 | 2.4 | 2.5 | 2.1 |
| 7 | 2.9 | 1.9 | 1.9 | 1.6 | 2.4 | 1.4 |
| 8 | 3.5 | 2.5 | 2.7 | 1.7 | 3.1 | 1.4 |
| Total | 2.9 | 1.4 | 2.4 | 1.2 | 2.7 | 1.0 |
| Current cigar use | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 5.8 | 4.4 | 2.9 | 3.1 | 4.5 | 3.3 |
| Black | 11.5 | 2.7 | 8.4 | 2.6 | 9.9 | 1.8 |
| Hispanic | 7.5 | 3.5 | 6.7 | 4.1 | 7.3 | 3.0 |
| Other | _ | _ | _ | _ | 5.8 | 4.0 |
| Grade | | | | | | |
| 6 | 7.8 | 4.8 | 4.5 | 3.5 | 6.2 | 3.0 |
| 7 | 12.2 | 4.1 | 7.5 | 3.6 | 9.8 | 2.9 |
| 8 | 9.3 | 2.7 | 9.9 | 2.6 | 9.6 | 1.6 |
| Total | 9.8 | 1.9 | 7.3 | 2.1 | 8.6 | 1.5 |
| Current tobacco use | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 10.0 | 6.5 | 14.4 | 6.0 | 11.9 | 4.5 |
| Black | 12.6 | 2.4 | 11.1 | 2.9 | 11.9 | 1.8 |
| Hispanic | 11.1 | 4.6 | 14.9 | 5.5 | 13.2 | 3.6 |
| Other | _ | _ | _ | _ | 12.1 | 5.4 |
| Grade | | | | | | |
| 6 | 7.9 | 4.3 | 8.6 | 4.6 | 8.2 | 2.6 |
| 7 | 13.2 | 3.8 | 11.5 | 4.2 | 12.3 | 2.8 |
| 8 | 14.5 | 4.4 | 16.1 | 3.8 | 15.2 | 2.5 |
| Total | 11.8 | 2.1 | 12.2 | 2.2 | 12.1 | 1.5 |

^{*}Used chewing tobacco, snuff, or dip on ≥ 1 of the 30 days preceding the survey.

 $^{^{\}dagger}$ Smoked cigars, cigarillos, or little cigars on ≥ 1 of the 30 days preceding the survey.

[§]Reported current cigarette use, current cigar use, or had used chewing tobacco, snuff, or dip on ≥ 1 of the 30 days preceding the survey.

^{195%} confidence interval.

^{**}Non-Hispanic.

^{††}Not available.

Table 9k. Percentage of Middle School Students Who Had Ever Had a Drink of Alcohol,* Who Had Ever Used Marijuana, and Who Had Ever Used Cocaine,† by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | Male | | Female | | Total | |
|-----------------------|------|---------|--------|--------|-------|--------|
| Question | % | CI§ (±) | % | CI (±) | % | CI (±) |
| Lifetime alcohol use | | | | | | |
| Race/Ethnicity | | | | | | |
| White ¹ | 48.9 | 9.6 | 49.6 | 9.1 | 49.2 | 6.8 |
| Black ¹ | 45.6 | 6.6 | 51.2 | 4.7 | 48.3 | 3.9 |
| Hispanic | 44.9 | 10.2 | 51.8 | 9.3 | 48.3 | 7.5 |
| Other | ** | _ | _ | _ | 50.0 | 7.4 |
| Grade | | | | | | |
| 6 | 46.2 | 10.7 | 42.1 | 8.0 | 44.1 | 6.7 |
| 7 | 38.6 | 7.6 | 52.8 | 5.3 | 45.4 | 4.7 |
| 8 | 54.7 | 5.1 | 57.4 | 6.7 | 56.0 | 4.1 |
| Total | 46.2 | 5.0 | 51.1 | 3.9 | 48.6 | 3.3 |
| ifetime marijuana use | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 12.1 | 7.2 | 8.4 | 6.7 | 10.5 | 5.9 |
| Black | 30.0 | 5.0 | 18.8 | 3.9 | 24.2 | 3.8 |
| Hispanic | 22.8 | 10.2 | 16.8 | 8.0 | 19.8 | 6.9 |
| Other | _ | _ | _ | _ | 17.9 | 6.7 |
| Grade | | | | | | |
| 6 | 17.3 | 6.9 | 6.1 | 4.7 | 11.8 | 4.6 |
| 7 | 21.6 | 5.9 | 15.1 | 4.8 | 18.2 | 4.1 |
| 8 | 36.5 | 6.5 | 29.5 | 5.2 | 33.0 | 4.5 |
| Total | 25.0 | 4.6 | 17.1 | 3.5 | 21.1 | 3.4 |
| ifetime cocaine use | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 2.2 | 2.6 | 2.8 | 2.9 | 2.4 | 2.2 |
| Black | 4.2 | 2.3 | 2.4 | 1.0 | 3.3 | 1.3 |
| Hispanic | 5.4 | 3.9 | 4.2 | 4.0 | 4.8 | 2.6 |
| Other | _ | _ | _ | _ | 5.4 | 3.7 |
| Grade | | | | | | |
| 6 | 4.1 | 3.5 | 2.2 | 2.1 | 3.2 | 2.3 |
| 7 | 3.6 | 2.4 | 3.4 | 2.3 | 3.5 | 1.7 |
| 8 | 4.5 | 2.7 | 2.9 | 2.0 | 3.7 | 1.6 |
| Total | 4.3 | 1.8 | 2.8 | 1.1 | 3.6 | 1.1 |

^{*}Ever had a drink of alcohol, other than a few sips.

 $^{^{\}dagger}\textsc{Ever}$ used any form of cocaine (e.g., powder, "crack," or "freebase").

^{§95%} confidence interval.

¹Non-Hispanic.

^{**}Not available.

Table 10k. Percentage of Middle School Students Who Had Ever Used Inhalants* and Who Had Ever Used Steroids, by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | N | 1ale | Female | | Total | |
|-----------------------|-----|---------------------|--------|--------|-------|--------|
| Question | % | CI [†] (±) | % | CI (±) | % | CI (±) |
| Lifetime inhalant use | | | | | | |
| Race/Ethnicity | | | | | | |
| White [§] | 6.4 | 4.2 | 12.5 | 5.5 | 9.1 | 3.5 |
| Black [§] | 5.3 | 2.3 | 5.5 | 2.0 | 5.4 | 1.6 |
| Hispanic | 8.4 | 6.4 | 9.8 | 5.2 | 9.1 | 5.6 |
| Other | 1 | | _ | | 10.5 | 4.7 |
| Grade | | | | | | |
| 6 | 7.1 | 4.5 | 9.7 | 4.6 | 8.4 | 3.7 |
| 7 | 6.5 | 3.2 | 4.7 | 2.4 | 5.6 | 2.3 |
| 8 | 5.2 | 2.3 | 7.7 | 3.5 | 6.5 | 2.0 |
| Total | 6.3 | 1.9 | 7.3 | 2.0 | 6.8 | 1.6 |
| Lifetime steroid use | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 0.6 | 1.1 | 2.8 | 3.5 | 1.5 | 1.8 |
| Black | 2.6 | 1.8 | 1.7 | 1.3 | 2.2 | 0.9 |
| Hispanic | 1.9 | 2.0 | 2.2 | 2.2 | 2.1 | 1.2 |
| Other | | | _ | | 4.7 | 3.4 |
| Grade | | | | | | |
| 6 | 2.8 | 3.5 | 3.0 | 2.7 | 2.9 | 2.1 |
| 7 | 2.0 | 1.7 | 2.2 | 1.6 | 2.1 | 1.1 |
| 8 | 2.4 | 1.7 | 0.8 | 0.9 | 1.6 | 1.0 |
| Total | 2.4 | 1.3 | 2.0 | 1.1 | 2.2 | 0.8 |

^{*}Sniffed glue, breathed the contents of spray cans, or inhaled any paints or sprays to get high.

^{†95%} confidence interval.

[§]Non-Hispanic.

¹Not available.

Table I Ik. Percentage of Middle School Students Who Had Initiated Drug-Related Behaviors Before Age I I Years, by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | | M ale | | male | Total | |
|----------------------------|----------------------------------|--------------|------|--------|-------|--------|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) |
| Smoked a whole cigarett | e before age 11 years | s | | | | |
| Race/Ethnicity | | | | | | |
| White [†] | 13.7 | 8.2 | 8.2 | 4.5 | 11.3 | 4.8 |
| Black [†] | 8.2 | 3.1 | 9.4 | 2.8 | 8.8 | 2.2 |
| Hispanic | 8.1 | 4.1 | 11.2 | 5.1 | 9.7 | 3.4 |
| Other | § | _ | _ | _ | 9.0 | 4.6 |
| Grade | | | | | | |
| 6 | 8.3 | 4.3 | 6.7 | 4.2 | 7.5 | 3.2 |
| 7 | 8.5 | 3.4 | 10.4 | 3.1 | 9.4 | 2.6 |
| 8 | 11.3 | 4.2 | 10.4 | 3.3 | 10.9 | 2.6 |
| Total | 9.4 | 2.4 | 9.2 | 2.0 | 9.3 | 1.6 |
| Had first drink of alcohol | before age II years ¹ | | | | | |
| Race/Ethnicity | | | | | | |
| White | 23.3 | 9.1 | 15.1 | 7.9 | 19.7 | 7.2 |
| Black | 17.9 | 4.8 | 18.6 | 5.7 | 18.2 | 3.9 |
| Hispanic | 20.9 | 7.6 | 18.0 | 6.8 | 19.7 | 5.2 |
| Other | _ | _ | _ | _ | 17.5 | 5.8 |
| Grade | | | | | | |
| 6 | 24.0 | 6.8 | 21.6 | 9.5 | 22.8 | 5.7 |
| 7 | 14.6 | 5.6 | 18.6 | 7.3 | 16.7 | 5.2 |
| 8 | 20.5 | 4.9 | 13.2 | 4.3 | 16.8 | 3.4 |
| Total | 19.6 | 3.6 | 17.7 | 4.3 | 18.7 | 3.0 |
| Tried marijuana before aş | ge II years | | | | | |
| Race/Ethnicity | - | | | | | |
| White | 0.0 | 0.0 | 0.8 | 1.5 | 0.3 | 0.7 |
| Black | 4.1 | 1.6 | 1.9 | 1.2 | 3.0 | 1.0 |
| Hispanic | 4.3 | 3.2 | 5.1 | 3.4 | 4.9 | 2.7 |
| Other | _ | _ | _ | _ | 3.3 | 2.5 |
| Grade | | | | | | |
| 6 | 2.6 | 2.5 | 0.9 | 1.2 | 1.8 | 1.5 |
| 7 | 3.8 | 2.4 | 3.1 | 1.8 | 3.4 | 1.7 |
| 8 | 4.0 | 1.9 | 2.4 | 1.6 | 3.2 | 1.1 |
| Total | 3.5 | 1.2 | 2.2 | 1.0 | 3.0 | 0.8 |

^{*95%} confidence interval.

[†]Non-Hispanic.

[§]Not available.

¹Other than a few sips.

Table 12k. Percentage of Middle School Students Who Had Engaged in Sexual Behaviors, by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | Male | | Female | | Total | |
|----------------------------------|-----------------|---------|--------|--------|-------|--------|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) |
| Ever had sexual intercourse | | | | | | |
| Race/Ethnicity | | | | | | |
| White [†] | 19.1 | 9.2 | § | _ | 15.8 | 6.1 |
| Black [†] | 51.8 | 6.9 | 19.7 | 5.5 | 35.2 | 4.9 |
| Hispanic | 18.3 | 10.3 | 14.6 | 7.1 | 16.3 | 6.8 |
| Other | _ | _ | _ | _ | 24.3 | 7.3 |
| Grade | | | | | | |
| 6 | 25.0 | 11.9 | 6.9 | 4.6 | 15.8 | 5.9 |
| 7 | 40.0 | 8.1 | 18.3 | 4.7 | 29.2 | 4.7 |
| 8 | 54.1 | 7.1 | 27.3 | 6.6 | 40.5 | 5.7 |
| Total | 39.8 | 6.2 | 17.8 | 4.0 | 28.8 | 4.1 |
| Had first sexual intercourse bef | fore age II yea | ırs | | | | |
| Race/Ethnicity | | | | | | |
| White | 9.8 | 7.7 | _ | _ | 7.2 | 4.3 |
| Black | 19.8 | 4.5 | 2.9 | 1.6 | 11.1 | 2.5 |
| Hispanic | 6.5 | 4.9 | 2.9 | 2.6 | 4.6 | 2.7 |
| Other | _ | _ | _ | _ | 4.3 | 3.1 |
| Grade | | | | | | |
| 6 | 15.1 | 9.4 | 2.4 | 2.4 | 8.7 | 4.5 |
| 7 | 16.0 | 5.8 | 5.1 | 2.3 | 10.6 | 3.3 |
| 8 | 14.7 | 4.5 | 1.1 | 1.2 | 7.8 | 2.1 |
| Total | 15.2 | 3.5 | 2.9 | 1.3 | 9.1 | 1.8 |
| Had ≥ 3 sex partners | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 4.7 | 4.3 | _ | _ | 3.4 | 2.4 |
| Black | 24.1 | 5.7 | 4.4 | 2.0 | 13.9 | 3.1 |
| Hispanic | 5.4 | 2.8 | 4.5 | 3.9 | 4.9 | 2.5 |
| Other | _ | _ | _ | _ | 9.3 | 3.8 |
| Grade | | | | | | |
| 6 | 11.5 | 7.5 | 2.0 | 2.0 | 6.6 | 4.2 |
| 7 | 16.5 | 5.5 | 3.9 | 2.0 | 10.2 | 3.0 |
| 8 | 23.6 | 6.0 | 6.0 | 3.9 | 14.8 | 3.6 |
| Total | 17.3 | 4.2 | 4.0 | 1.6 | 10.6 | 2.4 |

^{*95%} confidence interval.

[†]Non-Hispanic.

[§]Not available.

Table 13k. Percentage of Middle School Students Who Had Used a Condom During Last Sexual Intercourse* and Who Had Ever Been Taught About Acquired Immunodeficiency Syndrome (AIDS) or Human Immunodeficiency Virus (HIV) Infection in School, by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | <u> </u> | Male | | Female | | otal . |
|---------------------------|--------------------|---------------------|------|--------|------|--------|
| Question | % | CI [†] (±) | % | CI (±) | % | CI (±) |
| Used condom during last | sexual intercourse | | | | | |
| Race/Ethnicity | | | | | | |
| White§ | 1 | _ | _ | _ | _ | _ |
| Black [§] | 79.8 | 7.0 | _ | _ | 77.3 | 6.2 |
| Hispanic | _ | _ | _ | _ | _ | _ |
| Other | _ | _ | _ | _ | _ | _ |
| Grade | | | | | | |
| 6 | _ | _ | _ | _ | _ | _ |
| 7 | _ | _ | _ | _ | 76.7 | 7.1 |
| 8 | 82.8 | 5.4 | _ | _ | 81.7 | 5.4 |
| Total | 78.1 | 7.3 | 71.8 | 9.6 | 76.2 | 5.9 |
| Had been taught in school | about AIDS or HIV | infection | | | | |
| Race/Ethnicity | | | | | | |
| White | 63.1 | 15.2 | 67.9 | 11.3 | 65.2 | 12.3 |
| Black | 74.5 | 5.2 | 74.4 | 5.7 | 74.4 | 4.3 |
| Hispanic | 54.5 | 11.7 | 56.1 | 14.6 | 55.3 | 11.7 |
| Other | _ | _ | _ | _ | 67.6 | 10.0 |
| Grade | | | | | | |
| 6 | 48.4 | 9.5 | 58.6 | 9.5 | 53.3 | 8.5 |
| 7 | 76.5 | 6.6 | 70.0 | 8.2 | 73.3 | 6.3 |
| 8 | 80.3 | 5.4 | 82.2 | 4.4 | 81.2 | 4.0 |
| Total | 68.7 | 5.4 | 70.4 | 5.2 | 69.5 | 4.7 |

^{*}Among the 28.8% of students who ever had sexual intercourse.

^{†95%} confidence interval.

[§]Non-Hispanic.

¹Not available.

Table 14k. Percentage of Middle School Students Who Had Participated in Sufficient Vigorous Physical Activity,* Who Had Played on Any Sports Teams,† and Who Had Watched ≥ 3 Hours per Day of TV,§ by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | Male | | Female | | Total | |
|-----------------------------------|------------------|---------------------|--------|--------|-------|--------|
| Question | % | CI ¹ (±) | % | CI (±) | % | CI (±) |
| Participated in sufficient vigoro | us physical acti | vity | | | | |
| Race/Ethnicity | | | | | | |
| White** | 81.2 | 6.7 | 71.1 | 10.0 | 76.9 | 5.1 |
| Black** | 59.6 | 5.5 | 53.3 | 6. l | 56.5 | 4.1 |
| Hispanic | 71.4 | 8.4 | 62.0 | 6.7 | 66.8 | 5.4 |
| Other | | _ | _ | _ | 58.9 | 7.5 |
| Grade | | | | | | |
| 6 | 60.6 | 9.3 | 60.1 | 9.4 | 60.3 | 6.4 |
| 7 | 62.8 | 7.2 | 55.6 | 7.2 | 59.5 | 6.4 |
| 8 | 70.9 | 5.5 | 56.8 | 6.7 | 64.0 | 4.6 |
| Total | 64.5 | 4.5 | 57.6 | 4.3 | 61.2 | 3.1 |
| Played on any sports teams | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 51.5 | 9.3 | 46.2 | 14.0 | 49.2 | 7.5 |
| Black | 59.8 | 5.7 | 42.4 | 4.2 | 50.9 | 3.4 |
| Hispanic | 52.4 | 11.8 | 40.5 | 7.6 | 46.6 | 7.0 |
| Other | _ | _ | _ | _ | 48.9 | 8.6 |
| Grade | | | | | | |
| 6 | 56.3 | 9.3 | 45.6 | 8.5 | 50.9 | 7.0 |
| 7 | 56.9 | 5.3 | 41.5 | 5.6 | 49.5 | 4.0 |
| 8 | 57.2 | 7.3 | 41.0 | 5.8 | 49.2 | 4.4 |
| Total | 56.8 | 4.5 | 42.7 | 3.9 | 49.9 | 3.0 |
| Watched \geq 3 hours per day of | ΓV | | | | | |
| Race/Ethnicity | | | | | | |
| White | 54.7 | 10.0 | 44.5 | 8.3 | 50.3 | 7.0 |
| Black | 69.6 | 5.0 | 69.2 | 4.8 | 69.5 | 3.5 |
| Hispanic | 45.3 | 8.9 | 51.2 | 10.0 | 48.4 | 7.3 |
| Other | _ | _ | _ | _ | 58.1 | 7.1 |
| Grade | | | | | | |
| 6 | 55.6 | 9.6 | 56.1 | 10.2 | 55.9 | 7.7 |
| 7 | 64.7 | 6.6 | 66.4 | 5.6 | 65.7 | 4.4 |
| 8 | 66.9 | 6.0 | 63.7 | 4.3 | 65.3 | 3.8 |
| Total | 62.5 | 3.7 | 62.2 | 4.0 | 62.5 | 2.8 |

^{*}Exercised or participated in physical activities that made them sweat and breathe hard for ≥ 20 minutes on ≥ 3 of the 7 days preceding the survey (e.g., basketball, soccer, running, swimming laps, fast bicycling, fast dancing, or similar aerobic activities).

[†]Run by their school or community groups.

[§]On an average school day.

^{195%} confidence interval.

^{**}Non-Hispanic.

^{††}Not available.

Table 15k. Percentage of Middle School Students Who Were Enrolled in Physical Education (PE) Class* and Attended PE Class Daily,† by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| Question | Male | | Female | | Total | |
|---------------------------|------|---------|--------|--------|-------|--------|
| | % | CI§ (±) | % | CI (±) | % | CI (±) |
| Were enrolled in PE class | | | | | | |
| Race/Ethnicity | | | | | | |
| White ¹ | 81.5 | 8.7 | 86.2 | 6.7 | 83.6 | 6.4 |
| Black ¹¹ | 73.0 | 6.3 | 78.0 | 5.9 | 75.5 | 4.7 |
| Hispanic | 77.8 | 9.0 | 75.7 | 9.2 | 76.8 | 8.2 |
| Other | ** | | _ | _ | 72.2 | 7.4 |
| Grade | | | | | | |
| 6 | 73.9 | 10.8 | 80.0 | 7.0 | 77.0 | 7.9 |
| 7 | 74.7 | 7.7 | 72.4 | 8.1 | 73.6 | 6.1 |
| 8 | 76.0 | 6.8 | 83.4 | 6.6 | 79.6 | 5.7 |
| Total | 74.7 | 4.6 | 78.5 | 4.5 | 76.6 | 3.5 |
| Attended PE class daily | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 26.2 | 10.2 | 20.4 | 9.7 | 23.6 | 8.9 |
| Black | 28.1 | 7.1 | 31.1 | 7.8 | 29.6 | 6.4 |
| Hispanic | 27.3 | 11.5 | 21.3 | 10.6 | 24.3 | 9.8 |
| Other | _ | _ | _ | _ | 24.0 | 8.0 |
| Grade | | | | | | |
| 6 | 24.7 | 16.1 | 24.1 | 11.8 | 24.4 | 12.7 |
| 7 | 25.1 | 11.6 | 23.4 | 9.8 | 24.2 | 9.8 |
| 8 | 31.2 | 8.4 | 36.7 | 15.5 | 33.9 | 11.2 |
| Total | 26.9 | 5.7 | 28.2 | 5.7 | 27.5 | 4.8 |

^{*}On one or more days in an average week when they were in school.

 $^{^{\}dagger}5$ days in an average week when they were in school.

^{§95%} confidence interval.

¹Non-Hispanic.

^{**}Not available.

Table 16k. Percentage of Middle School Students Who Were at Risk for Becoming* or Were Overweight,† by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | Male | | Female | | Total | |
|---------------------------------|------|---------------------|--------|--------|-------|--------|
| Question | % | CI [§] (±) | % | CI (±) | % | CI (±) |
| At risk for becoming overweight | | | | | | |
| Race/Ethnicity | | | | | | |
| White ¹ | 20.0 | 9.2 | ** | _ | 19.2 | 5.9 |
| Black ¹ | 17.1 | 5.2 | 24.9 | 4.2 | 21.0 | 3.6 |
| Hispanic | 23.3 | 6.5 | 23.6 | 8.0 | 23.4 | 4.6 |
| Other | _ | _ | _ | _ | 15.4 | 6.2 |
| Grade | | | | | | |
| 6 | 22.3 | 6.0 | 17.0 | 6.9 | 19.9 | 5.0 |
| 7 | 18.7 | 6.6 | 23.3 | 5.4 | 21.0 | 4.2 |
| 8 | 15.8 | 4.7 | 27.2 | 5.3 | 21.5 | 3.5 |
| Total | 18.6 | 3.4 | 23.3 | 3.5 | 20.9 | 2.5 |
| Overweight | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 18.3 | 5.8 | _ | _ | 12.7 | 4.0 |
| Black | 16.1 | 4.1 | 15.2 | 4.2 | 15.6 | 3.0 |
| Hispanic | 20.5 | 8.9 | 18.5 | 7.6 | 19.5 | 6.3 |
| Other | _ | _ | _ | _ | 15.8 | 7.6 |
| Grade | | | | | | |
| 6 | 21.4 | 7.5 | 15.7 | 8.5 | 18.8 | 6.5 |
| 7 | 15.8 | 4.3 | 15.3 | 5.6 | 15.6 | 3.4 |
| 8 | 15.9 | 3.2 | 13.1 | 4.1 | 14.5 | 2.7 |
| Total | 17.5 | 3.0 | 14.5 | 3.3 | 16.0 | 2.4 |

^{*}Students who were \geq 85th percentile but < 95th percentile for body mass index, by age and sex, based on reference data.

 $^{^{\}dagger}$ Students who were \geq 95th percentile for body mass index, by age and sex, based on reference data.

^{§95%} confidence interval.

¹Non-Hispanic.

^{**}Not available.

Table 17k. Percentage of Middle School Students Who Described Themselves as Slightly or Very Overweight and Who Were Trying To Lose Weight, by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | Male | | Female | | Total | |
|------------------------------------|------|---------|--------|--------|-------|--------|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) |
| Described themselves as overweight | | | | | | |
| Race/Ethnicity | | | | | | |
| White [†] | 26.0 | 9.9 | 30.4 | 7.8 | 27.8 | 7.4 |
| Black [†] | 14.7 | 3.3 | 15.8 | 3.4 | 15.2 | 2.5 |
| Hispanic | 25.1 | 5.7 | 31.8 | 10.3 | 28.5 | 5.2 |
| Other | § | _ | _ | | 19.5 | 5.7 |
| Grade | | | | | | |
| 6 | 21.4 | 5.5 | 16.5 | 5.2 | 19.0 | 4.2 |
| 7 | 17.1 | 5.6 | 19.9 | 4.8 | 18.3 | 4.2 |
| 8 | 17.4 | 3.2 | 25.1 | 5.5 | 21.2 | 3.2 |
| Total | 18.4 | 2.8 | 20.6 | 2.9 | 19.5 | 2.1 |
| Were trying to lose weight | | | | | | |
| Race/Ethnicity | | | | | | |
| White | 45.4 | 8.4 | 49.8 | 13.9 | 47.3 | 8.5 |
| Black | 33.2 | 5.2 | 39.9 | 4.4 | 36.6 | 3.2 |
| Hispanic | 41.6 | 8.3 | 54.7 | 9.7 | 48.0 | 7.1 |
| Other | _ | _ | _ | _ | 40.1 | 6.4 |
| Grade | | | | | | |
| 6 | 40.5 | 7.4 | 41.8 | 7.3 | 41.1 | 5.8 |
| 7 | 36.6 | 5.6 | 40.2 | 4.4 | 38.2 | 3.1 |
| 8 | 33.0 | 4.0 | 49.6 | 6.6 | 41.1 | 4.0 |
| Total | 36.7 | 3.6 | 43.8 | 3.6 | 40.1 | 2.5 |

^{*95%} confidence interval.

[†]Non-Hispanic.

[§]Not available.

Table 18k. Percentage of Middle School Students Who Had Ever Engaged in Healthy Behaviors Associated with Weight Control To Lose Weight or To Keep from Gaining Weight, by Sex, Race/Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| • | • | • | | | | |
|---|--------------------------|-------------------|----------------|-----------------|--------------|--------|
| Question | | Male | | Female | | otal |
| | % | CI* (±) | % | CI (±) | % | CI (±) |
| Ate less food, fewer calor | ries, or foods low in fa | at to lose weight | or to keep fro | om gaining weig | ht | |
| Race/Ethnicity | | | | | | |
| White [†] | 37.2 | 9.5 | 60.5 | 10.9 | 47.3 | 6.6 |
| Black [†] | 33.7 | 5.6 | 37. I | 3.9 | 35.6 | 3.1 |
| Hispanic | 45.8 | 8.9 | 52.6 | 7.5 | 49.2 | 5.1 |
| Other | § | _ | _ | _ | 45. I | 7.1 |
| Grade | | | | | | |
| 6 | 47.7 | 8.8 | 43.4 | 7.3 | 45.6 | 6.1 |
| 7 | 30.1 | 6.8 | 39.4 | 5.0 | 34.9 | 4.0 |
| 8 | 32.8 | 4.9 | 47.2 | 5.9 | 39.9 | 4.0 |
| Total | 36.8 | 4.5 | 43.3 | 3.2 | 40.1 | 2.9 |
| Exercised to lose weight | or to keep from gain | ing weight | | | | |
| Race/Ethnicity | | | | | | |
| White | 61.1 | 9.0 | 70. I | 9.2 | 65.0 | 5.8 |
| Black | 60.9 | 5.2 | 57.0 | 5.6 | 59.0 | 3.6 |
| Hispanic | 63.9 | 7.5 | 70.5 | 7.0 | 67.3 | 5.8 |
| Other | _ | _ | _ | _ | 63.9 | 6.9 |
| Grade | | | | | | |
| 6 | 62.0 | 8.1 | 60.2 | 8.7 | 61.1 | 6.4 |
| 7 | 57.8 | 7.0 | 60. I | 6.3 | 59.1 | 4.5 |
| 8 | 64.9 | 4.1 | 63.4 | 5.6 | 64.2 | 4.1 |
| Total | 61.4 | 3.7 | 61.3 | 4.2 | 61.5 | 2.9 |

^{*95%} confidence interval.

[†]Non-Hispanic.

[§]Not available.

Table 19k. Percentage of Middle School Students Who Had Ever Engaged in Unhealthy Behaviors Associated with Weight Control To Lose Weight or To Keep from Gaining Weight, by Sex, Race/ Ethnicity, and Grade—Milwaukee PS, WI, Middle School Youth Risk Behavior Survey, 2003

| | | Male | | Female | | Total | |
|---------------------------|-------------------------|-------------------|--------------|--------------------|------|--------|--|
| Question | % | CI* (±) | % | CI (±) | % | CI (±) | |
| Went without eating for | ≥ 24 hours to lose w | eight or to keep | from gaining | weight | | - | |
| Race/Ethnicity | | | | | | | |
| White [†] | 16.3 | 5.7 | 27.8 | 7.4 | 21.2 | 4.5 | |
| Black [†] | 20.9 | 4.2 | 25.6 | 3.9 | 23.2 | 3.3 | |
| Hispanic | 17.1 | 6.2 | 23.7 | 6.3 | 20.4 | 5.1 | |
| Other | § | _ | _ | _ | 20.2 | 7.4 | |
| Grade | | | | | | | |
| 6 | 20.4 | 5.8 | 25.7 | 5.5 | 23.0 | 4.7 | |
| 7 | 17.1 | 4.5 | 20.3 | 4.4 | 18.6 | 3.0 | |
| 8 | 21.0 | 4.5 | 29.8 | 5.4 | 25.3 | 3.9 | |
| Total | 19.3 | 3.0 | 25.3 | 3.0 | 22.2 | 2.5 | |
| Took diet pills, powders, | or liquids to lose wei | ght or to keep fi | om gaining w | eight ¹ | | | |
| Race/Ethnicity | | | | | | | |
| White | 1.0 | 1.5 | 8.4 | 5.5 | 4.2 | 2.5 | |
| Black | 6.4 | 2.7 | 6.6 | 2.6 | 6.4 | 2.0 | |
| Hispanic | 9.3 | 4.2 | 8.5 | 4.1 | 8.9 | 3.5 | |
| Other | _ | _ | _ | _ | 4.4 | 3.3 | |
| Grade | | | | | | | |
| 6 | 6.2 | 3.6 | 3.4 | 2.6 | 4.8 | 2.3 | |
| 7 | 4.3 | 2.7 | 7.7 | 4.3 | 5.9 | 2.8 | |
| 8 | 6.7 | 3.4 | 10.2 | 2.9 | 8.4 | 2.3 | |
| Total | 5.7 | 2.0 | 7.2 | 2.1 | 6.5 | 1.5 | |
| omited or took laxative | es to lose weight or to | keep from gain | ing weight | | | | |
| Race/Ethnicity | | | | | | | |
| White | 0.5 | 1.0 | 9.3 | 6.4 | 4.3 | 2.8 | |
| Black | 4.7 | 2.1 | 5.4 | 1.8 | 5.0 | 1.1 | |
| Hispanic | 7.3 | 3.8 | 10.0 | 4.9 | 8.6 | 3.2 | |
| Other | _ | _ | _ | _ | 5.3 | 4.1 | |
| Grade | | | | | | | |
| 6 | 5.3 | 3.3 | 7.8 | 3.2 | 6.6 | 2.2 | |
| 7 | 2.7 | 2.0 | 5.7 | 2.2 | 4.1 | 1.5 | |
| 8 | 4.2 | 2.0 | 7.4 | 2.3 | 5.8 | 1.4 | |
| Total | 4.2 | 1.6 | 6.9 | 1.5 | 5.5 | 1.1 | |

^{*95%} confidence interval.

[†]Non-Hispanic.

[§]Not available.

[¶]Without a doctor's advice.